



Orexin receptor-1 expression in adult rodent neurogenic regions: evidence from the subgranular zone and the median eminence

Ekspresja receptora oreksynowego-1 w regionach neurogenezy dorosłych gryzoni: strefie podziarnistej oraz wyniosłości pośrodkowej

Antonello Pinna, Artur Pałasz 

Department of Histology, Faculty of Medical Sciences in Katowice, Medical University of Silesia, Katowice, Poland

ABSTRACT

INTRODUCTION: Orexin signaling plays a vital role in regulating autonomic and cognitive functions, including sleep-wake cycles, feeding, and memory. Orexin receptor-1 (OX₁R), a key component of this system, may also influence adult neurogenesis. This study examined OX₁R expression in both the classic (hippocampal) and non-classic (hypothalamic) neurogenic regions of the adult rodent brain.

MATERIAL AND METHODS: Adult rodent brains were fixed, paraffin-embedded, and sectioned coronally. Immunohistochemistry and immunofluorescence were performed using antibodies against OX₁R, DCX, and TUC-4, followed by fluorophore- or diaminobenzidine-based detection. Negative controls were included to ensure specificity.

RESULTS: OX₁R-positive cells were localized primarily to the subgranular zone (SGZ) of the dentate gyrus and β 2 tanycytes of the median eminence showed uniform OX₁R expression in both somata and vascular-directed processes. Morphological variation was observed between species, with diverse perikaryon shapes in mice and predominantly elongated, multipolar forms in rats.

CONCLUSIONS: This study revealed, for the first time, region-specific OX₁R expression in the SGZ in β 2 tanycytes of the median eminence. These findings suggest a potential role for orexin signaling in adult neurogenesis.

KEYWORDS

orexin, adult neurogenesis, hippocampus, tanycytes

Received: 14.05.2025

Revised: 11.06.2025

Accepted: 18.06.2025

Published online: 23.09.2025

Address for correspondence: prof. dr hab. n. med. Artur Pałasz, Zakład Histologii, Wydział Nauk Medycznych w Katowicach SUM, ul. Medyków 18, 40-752 Katowice, tel. +48 32 208 83 77, e-mail: apalasz@sum.edu.pl



This is an open access article made available under the terms of the Creative Commons Attribution-ShareAlike 4.0 International (CC BY-SA 4.0) license, which defines the rules for its use. It is allowed to copy, alter, distribute and present the work for any purpose, even commercially, provided that appropriate credit is given to the author and that the user indicates whether the publication has been modified, and when processing or creating based on the work, you must share your work under the same license as the original. The full terms of this license are available at <https://creativecommons.org/licenses/by-sa/4.0/legalcode>.

Publisher: Medical University of Silesia, Katowice, Poland



STRESZCZENIE

WSTĘP: Sygnalizacja oreksynergiczna odgrywa kluczową rolę w regulacji funkcji autonomicznych i poznawczych, w tym cyklu snu i czuwania oraz przyjmowania pokarmu. Receptor oreksyny-1 (*orexin receptor-1* – OX₁R), pierwszoplanowy element tego układu, może również wpływać na neurogenezę w mózgu dojrzałym. W przedstawionym badaniu zbadano ekspresję OX₁R zarówno w klasycznych (hipokamp), jak i nieklasycznych (podwzgórze) regionach neurogenicznych mózgu dorosłych gryzoni (szczurów i myszy).

MATERIAŁ I METODY: Mózgi dorosłych gryzoni utrwalono, zatopiono w parafinie i skrojono w płaszczyźnie poprzecznej. Badania immunohistochemiczne i fluorescencyjne wykonano z wykorzystaniem pierwszorzędowych przeciwciał selektywnych względem OX₁R, DCX i TUC-4, a następnie przeciwciał drugorzędowych sprzężonych z fluorochromami lub w reakcji diaminobenzydyny. W celu zapewnienia swoistości uwzględniono kontrole negatywne.

WYNIKI: Komórki OX₁R-pozytywne zlokalizowano głównie w strefie podziarnistej (*subgranular zone* – SGZ) zakrętu zębatego, β 2-tanycyty wyniosłości pośrodkowej wykazywały jednorodną ekspresję OX₁R zarówno w ich ciałach komórkowych, jak i w wypustkach okołonaczyniowych. Zaobserwowano istotne zróżnicowanie morfologiczne neuronów OX₁R-pozytywnych między badanymi gatunkami gryzoni – u myszy dominowały perykaryony różnokształtne, natomiast u szczurów komórki wydłużone i wielobiegunkowe.

WNIOSKI: Badanie to po raz pierwszy ujawniło specyficzną dla regionu ekspresję OX₁R w SGZ hipokampa oraz w β 2-tanycytach wyniosłości pośrodkowej podwzgórza. Wyniki sugerują potencjalną rolę sygnalizacji oreksynowej w regulacji neurogenety w mózgu dojrzałym.

SŁOWA KLUCZOWE

oreksyna, neurogeniza w mózgu dojrzałym, hipokamp, tanycyty

INTRODUCTION

Neurochemical research on brain neuropeptides involved in the regulation of adult neurogenesis has emerged as a highly dynamic area in contemporary neuroscience. Two well-established neurogenic niches have been identified in the vertebrate brain: the subgranular zone (SGZ) of the dentate gyrus and the subventricular zone (SVZ) beneath the ependyma of the lateral ventricles. The SGZ gives rise to granule neurons within the hippocampus, while the SVZ produces progenitor cells that migrate via the rostral migratory stream to the olfactory bulb, where they differentiate into specialized sensory interneurons. The SGZ niche is primarily composed of astrocytes and capillary endothelial cells, both of which are abundantly represented in this layer of the dentate gyrus [1,2]. In addition to these canonical regions, two potential sites of adult neurogenesis have recently been found within the hypothalamus: the hypothalamic ventricular zone (HVZ), situated along the lateral walls of the third ventricle, and the hypothalamic proliferative zone (HPZ), composed primarily of median eminence tanycytes [3,4] whose somata are situated along the floor of the third ventricle. Based on their anatomical distribution and the expression of specific markers related to lineage and differentiation, radial glia-like tanycytes are classified into four subtypes: α 1, α 2, β 1, and β 2 [5]. α 1 tanycytes are found at the ventromedial and dorsomedial nuclei of the hypothalamus, α 2 tanycytes are in the vicinity of the arcuate nuclei, and the β 1 and β 2 types are situated along the HVZ (ventral tanycytic zone) of the third ventricle [6], but extend their processes into the median eminence, in the tuberal region of the hypothalamus

[7]. Notably, β 2 tanycytes are characterized by their remarkable sensitivity to circulating hormones, metabolic regulators, and nutritional signals. Their somata, embedded in the ependymal lining, are optimally placed to detect molecular cues from the cerebrospinal fluid (CSF), while their elongated processes extend toward the median eminence, where they may interact with blood-borne signals [8]. Given their strategic location at the interface between the CSF and hypothalamic vasculature, β 2 tanycytes could be well positioned to respond to neuropeptides such as orexins. Tanycytes comprising the HVZ – including β 2 tanycytes in both rats and humans – express several neural precursor markers, including nestin [9], Sox2 [10], vimentin [11], and doublecortin-like protein [12], suggesting a role in adult neurogenesis.

In parallel, orexins exert broad regulatory effects across the central nervous system, influencing diverse physiological domains from sleep and homeostasis to memory, emotions, and reward [13,14]. Orexinergic perikarya, which co-express glutamate and dynorphine, are localized exclusively in the perifornical area and the lateral and posterior hypothalamus [15,16,17]. Their axons target multiple brain regions, including key neurogenic areas such as the hippocampus and the olfactory bulb [18,19]. Orexins act via two G-coupled receptors, known as OX₁R (orexin receptor-1) and OX₂R, showing significant homology among mammalian species [20]. OXRs have been identified in various brain regions, including the prefrontal and limbic cortices, the hippocampus, the amygdala, the bed nucleus of stria terminalis, and several hypothalamic and brainstem nuclei [21,22].

Building on recent evidence which suggests that certain hypothalamic orexin neurons project to neurogenic regions in the rat brain [23], we hypothesized that cells



within both the SGZ and the hypothalamic subependymal zone – particularly $\beta 2$ tanycytes – may be responsive to orexins via OX_1R signaling. This hypothesis is supported by the anatomical and functional profile of $\beta 2$ tanycytes: they are strategically positioned to sense hormonal and nutritional cues from both the CSF and the blood and they express multiple markers of neural precursors. These characteristics make them compelling candidates for mediating the neurogenic effects of orexins in the hypothalamus.

Thus, if $\beta 2$ tanycytes express OX_1R , they could serve as intermediaries through which orexins regulate adult neurogenesis, energy homeostasis, or hormonal feedback in hypothalamic circuits. To explore this possibility, we investigated OX_1R expression within both classic (hippocampal SGZ) and non-classic (hypothalamic median eminence) neurogenic zones.

MATERIAL AND METHODS

Animals

Male Sprague–Dawley rats and mice (N = 5 each) from the Medical University of Silesia Experimental Centre were housed at a temperature of 22°C with a regular 12/12 light/darkness cycle and access to standard Murigran chow and water *ad libitum*. The research was approved by the Local Ethical Commission for Animal Experimentation at the Medical University of Silesia (No. 36/2012) and all experimental procedures were conducted according to the NIH Guide for Care and Use of Animals.

Immunohistochemistry

The animals were quickly anesthetized with isoflurane inhalation and sacrificed. The brains were excised, fixed by immersion for 48 h in 4% paraformaldehyde PBS (pH 7.2–7.4) at 4°C, dehydrated via graded alcohols at room temperature, cleared in xylene, embedded in paraffin, and finally sectioned on a microtome (Leica Microsystems, Germany) in the coronal plane (–2.00 to –2.80 mm from the bregma) into 7- μ m thick slices, according to standard rat brain atlases.

For immunofluorescence, after blocking with 0.1% Triton X-100 (Sigma, T-7878) and 10% serum (horse normal serum for orexin, Vector Labs), sections were incubated overnight at 4°C with rabbit antibody against rat orexin-1 receptor (1:2000, Abcam). Additional

antibodies against adult neurogenesis markers were applied: goat anti-rat DCX (Santa Cruz Pharmaceuticals, 1:1000) and rabbit anti-rat TUC-4 (Abcam, 1:1000). Primary antibodies were followed by fluorochrome-conjugated secondary antibodies: goat anti-rabbit FITC (1:200, Abcam) for 1 h at 4°C. Finally, the sections were mounted on glass slides with DAPI-containing medium and coverslipped.

For diaminobenzidine (DAB) single staining, blocking (0.1% Triton X-100 and 10% serum) and administration of primary antibodies (1:1000) were followed by biotinylated anti-rabbit/anti-goat secondary antibodies for 30 min, and then an avidin-biotin-horseradish peroxidase complex (Vectastain ABC kit, Vector Labs) for another 30 min, before DAB was used to complete the reaction (1–2 min). The brain sections were dehydrated, mounted on glass slides with medium, and coverslipped. Sections incubated with rabbit/mouse IgG instead of primary antibody were used as negative controls in order to check the specificity of primary antibodies. All images were captured and analyzed with Nikon Coolpix optic systems and processed using Image ProPlus software (Media Cybernetics, USA). The same planes of the brain were chosen from each set of slides. On each section, we analyzed the morphology of immunopositive and immunonegative cells from the entire area of the dentate gyri and lateral hypothalamic regions.

RESULTS

Upon examining the hippocampal area of both mouse and rat brains, a distinct number of granular cells in the dentate gyrus exhibited OX_1R immunoreactivity. The highest aggregation of OX_1R -positive cells was observed almost exclusively within the SGZ, known as the classic site of adult neurogenesis. The OX_1R -immunoreactive cells in the mouse dentate gyri displayed a relatively wide spectrum of shapes: round, oval, fusiform, and even triangular (pyramidal) perikaryal, from which short dendrites sometimes emerged (Figure 1[A–F]). In the rat brain, oval, multipolar, and elongated OX_1R -immunoreactive somata were most common (Figure 1[G–H]). In the hypothalamus, the inner median eminence layer showed distinct, regular OX_1R expression. Columnar cell bodies and long processes running toward blood vessels indicated uniform, mild OX_1R immunoreactivity (Figure 2).




Fig. 1. Orexin receptor-1 (OX₁R)-expressing neurons in the hippocampus. Overview of neuroanatomical OX₁R immunodistribution in the mouse and rat Ammon's horn and dentate gyrus. Immunoperoxidase reaction with DAB staining in mouse hippocampus (A–D, F). Scale bars: 100 μ m (A–D), 25 μ m (F). Expression of adult neurogenesis markers doublecortin (DCX) and TUC-4 in the rat subventricular zone (E). Scale bars: 50 μ m (E), 25 μ m (insets). Immunofluorescence for OX₁R in the rat dentate gyrus (G–H). Scale bar: 50 μ m. gz – granular zone; ml – molecular layer; py – pyramidal layer; sgz – subgranular zone.




Fig. 2. Orexin receptor-1 (OX₁R)-immunopositive cells in the rat median eminence. Scale bars: 250 μ m (A–C), 50 μ m (A1–C1). EME – external medial eminence; IME – internal medial eminence; 3V – third ventricle.

DISCUSSION

Our study reveals a region-specific expression of OX₁R, in the SGZ of the dentate gyrus of the hippocampus in both rats and mice. For the first time, it reports strong OX₁R immunoreactivity in the majority of β 2 tanycytes within the median eminence of the hypothalamus.

Accumulating evidence suggests that the multidirectional nature of orexin signaling in the brain contributes to a wide range of autonomic and affective processes, including food intake, sleep-wake regulation, stress responses, reward and addiction mechanisms, and the

pathogenesis of various neuropsychiatric disorders [24,25,26,27,28,29]. The hippocampus is a well-established target of the orexinergic system, receiving dense projections from the lateral hypothalamus. Orexin receptors – OX₁R and OX₂R – are expressed in complementary patterns across the hippocampal subregions: OX₁R is found in CA1, CA2, and the dentate gyrus, while OX₂R is in the dentate gyrus and CA3 [21,22,30]. These receptors are implicated in diverse hippocampal functions, including reward processing, learning, memory [27,31,32,33], spatial cognition [34], stress response [35,36], and, notably, antinociception [37,38].



Orexin-A in particular has been shown to enhance hippocampal neurogenesis, potentially contributing to cognitive performance. However, whether adult neurogenesis influences the population of orexin-producing neurons remains uncertain [39,40]. For instance, local perfusion of orexin-A into the dentate gyrus promotes structural and functional plasticity [41], while transgenic mice lacking orexin neurons exhibit reduced hippocampal plasticity [42]. Intracerebroventricular orexin-A administration increases cellular proliferation in the dentate gyrus and alleviates depression-like symptoms [43,44]. Similarly, in pentylenetetrazol (PTZ)-kindled rat models of epilepsy, orexin-A boosts proliferation and promotes differentiation of neural precursors in the dentate gyrus [45]. The PTZ-induced kindling model is currently widely used in epileptogenic drug testing. As PTZ induces seizures via glutamatergic excitation and GABAergic inhibition [46], these findings suggest that orexin-A may counteract seizure-induced cognitive deficits through OX₁R-mediated neurogenesis. Reduced orexin-A levels following seizures may partly underlie such impairments [45]. Orexins also stimulate neurogenesis in the SVZ, although its functional relevance remains unclear [23]. It is still debated whether orexin signaling is essential for hippocampal neurogenesis and whether orexinergic neurons themselves can be replenished via adult neurogenesis [39,40].

Some evidence suggests that a limited population of newly generated hypothalamic neurons may express orexin [47]. In this brain region, adult neurogenesis is believed to support adaptive responses to dietary changes [48,49]. Tanycytes play a critical role, as they can sense hormonal and nutritional cues and can differentiate into orexigenic or anorexigenic neurons. However, the identity of tanycyte subtypes serving as true neural stem cells remains controversial. Notably, in seasonally breeding species, hypothalamic neurogenesis is modulated by day length, with increased levels being observed in the short photoperiod in sheep [50]. Similar changes in cell proliferation and tanycyte function align with feeding behavior and seasonal cues in hamsters [51,52].

Our study found that β 2 tanycytes in the inner layer of the median eminence displayed consistent OX₁R immunoreactivity in both their columnar somata and vascular-directed processes. Importantly, β 2 tanycytes are the most mitotically active tanycyte subtype [53]; they are especially neurogenic in younger animals [54]. These cells express high levels of Hes1 and Hes2 – markers of neural progenitors – and all newly generated neurons remain within the median eminence [54,55]. This local neurogenesis may support a stable population of sensory neurons responsible for monitoring CSF composition.

In addition, OX₁R-immunoreactive cells in dentate gyri displayed a variety of shapes. This intriguing diverse shape pattern is challenging to interpret and may reflect as yet unidentified interspecies differences in cellular morphology. Additionally, we cannot rule out the possibility that certain cells exhibit an irregular immunostaining pattern, potentially confined to the perikarya, thereby rendering some cellular processes undetectable.

It is important to acknowledge the limitations of our preliminary, predominantly qualitative morphological analysis. Future studies should aim to expand upon these findings by incorporating co-expression analyses of OX₁R with established markers of adult neurogenesis. While adult neurogenesis remains a prominent and debated topic in contemporary neuroscience, robust evidence supporting sustained, long-term neurogenesis in the adult human brain remains limited. Moreover, clinical approaches such as stem cell transplantation for neurodegenerative conditions, including Alzheimer's and Parkinson's diseases or ischemic brain injuries, have thus far yielded disappointing therapeutic outcomes [56]. In contrast, numerous animal studies continue to provide valuable insights into the mechanisms of adult neurogenesis, advancing our understanding of brain structure and function.

CONCLUSIONS

Our findings revealed a distinct expression pattern of OX₁R in the dentate gyrus of both rats and mice, specifically confined to the subgranular zone. Moreover, we demonstrated for the first time that the majority of β 2 tanycytes within the median eminence exhibit robust OX₁R immunoreactivity. These observations cautiously point to a previously underexplored link between orexin signaling and adult neurogenesis in both the classic hypothalamic neurogenic regions. Nonetheless, further research is necessary to elucidate the functional significance of these potential regulatory interactions.

Compliance with Ethical Standards

Ethical approval

All applicable international, national, and/or institutional guidelines for the care and use of animals were followed.

Conflict of interest

The authors declare no conflict of interest.



Authors' contribution

Study design – A. Pinna, A. Palasz

Data collection – A. Pinna

Data interpretation – A. Pinna, A. Palasz

Manuscript preparation – A. Pinna, A. Palasz

Literature research – A. Pinna

REFERENCES

1. Gonçalves J.T., Schafer S.T., Gage F.H. Adult neurogenesis in the hippocampus: from stem cells to behavior. *Cell* 2016; 167(4): 897–914, doi: 10.1016/j.cell.2016.10.021.
2. Rolando C., Taylor V. Neural stem cell of the hippocampus: development, physiology regulation, and dysfunction in disease. *Curr. Top Dev. Biol.* 2014; 107: 183–206, doi: 10.1016/B978-0-12-416022-4.00007-X.
3. Kokoeva M.V., Yin H., Flier J.S. Neurogenesis in the hypothalamus of adult mice: potential role in energy balance. *Science* 2005; 310(5748): 679–683, doi: 10.1126/science.1115360.
4. Pierce A.A., Xu A.W. De novo neurogenesis in adult hypothalamus as a compensatory mechanism to regulate energy balance. *J. Neurosci.* 2010; 30(2): 723–730, doi: 10.1523/JNEUROSCI.2479-09.2010.
5. Goodman T., Hajhosseini M.K. Hypothalamic tanyocytes-masters and servants of metabolic, neuroendocrine, and neurogenic functions. *Front Neurosci.* 2015; 9: 387, doi: 10.3389/fnins.2015.00387.
6. Mathew T.C. Regional analysis of the ependyma of the third ventricle of rat by light and electron microscopy. *Anat. Histol. Embryol.* 2008; 37(1): 9–18, doi: 10.1111/j.1439-0264.2007.00786.x.
7. Prevot V., Dehouck B., Sharif A., Ciofi P., Giacobini P., Clasadonte J. The versatile tanyocyte: a hypothalamic integrator of reproduction and energy metabolism. *Endocr. Rev.* 2018; 39(3): 333–368, doi: 10.1210/er.2017-00235.
8. Mullier A., Bouret S.G., Prevot V., Dehouck B. Differential distribution of tight junction proteins suggests a role for tanyocytes in blood-hypothalamus barrier regulation in the adult mouse brain. *J. Comp. Neurol.* 2010; 518(7): 943–962, doi: 10.1002/cne.22273.
9. Wei L.C., Shi M., Chen L.W., Cao R., Zhang P., Chan Y.S. Nestin-containing cells express glial fibrillary acidic protein in the proliferative regions of central nervous system of postnatal developing and adult mice. *Brain Res. Dev. Brain Res.* 2002; 139(1): 9–17, doi: 10.1016/S0165-3806(02)00509-6.
10. Lee D.A., Blackshaw S. Functional implications of hypothalamic neurogenesis in the adult mammalian brain. *Int. J. Dev. Neurosci.* 2012; 30(8): 615–621, doi: 10.1016/j.ijdevneu.2012.07.003.
11. Bolborea M., Dale N. Hypothalamic tanyocytes: potential roles in the control of feeding and energy balance. *Trends Neurosci.* 2013; 36(2): 91–100, doi: 10.1016/j.tins.2012.12.008.
12. Saaltink D.J., Håvik B., Verissimo C.S., Lucassen P.J., Vreugdenhil E. Doublecortin and doublecortin-like are expressed in overlapping and non-overlapping neuronal cell populations: implications for neurogenesis. *J. Comp. Neurol.* 2012; 520(13): 2805–2823, doi: 10.1002/cne.23144.
13. Bonnavion P., de Lecea L. Hypocretins in the control of sleep and wakefulness. *Curr. Neurol. Neurosci. Rep.* 2010; 10(3): 174–179, doi: 10.1007/s11910-010-0101-y.
14. Nattie E., Li A. Central chemoreception in wakefulness and sleep: evidence for a distributed network and a role for orexin. *J. Appl. Physiol.* (1985) 2010; 108(5): 1417–1424, doi: 10.1152/jappphysiol.01261.2009.
15. Peyron C., Tighe D.K., van den Pol A.N., de Lecea L., Heller H.C., Sutcliffe J.G. et al. Neurons containing hypocretin (orexin) project to multiple neuronal systems. *J. Neurosci.* 1998; 18(23): 9996–10015, doi: 10.1523/JNEUROSCI.18-23-09996.1998.
16. Torrealba F., Yanagisawa M., Saper C.B. Colocalization of orexin A and glutamate immunoreactivity in axon terminals in the tuberomammillary nucleus in rats. *Neuroscience* 2003; 119(4): 1033–1044, doi: 10.1016/S0306-4522(03)00238-0.
17. Chou T.C., Lee C.E., Lu J., Elmquist J.K., Hara J., Willie J.T. et al. Orexin (hypocretin) neurons contain dynorphin. *J. Neurosci.* 2001; 21(19): RC168, doi: 10.1523/JNEUROSCI.21-19-00003.2001.
18. Nambu T., Sakurai T., Mizukami K., Hosoya Y., Yanagisawa M., Goto K. Distribution of orexin neurons in the adult rat brain. *Brain Res.* 1999; 827(1–2): 243–260, doi: 10.1016/S0006-8993(99)01336-0.
19. van den Pol A.N. Hypothalamic hypocretin (orexin): robust innervation of the spinal cord. *J. Neurosci.* 1999; 19(8): 3171–3182, doi: 10.1523/JNEUROSCI.19-08-03171.1999.
20. Tsujino N., Sakurai T. Orexin/hypocretin: a neuropeptide at the interface of sleep, energy homeostasis, and reward system. *Pharmacol. Rev.* 2009; 61(2): 162–176, doi: 10.1124/pr.109.001321.
21. Marcus J.N., Aschkenasi C.J., Lee C.E., Chemelli R.M., Saper C.B., Yanagisawa M. et al. Differential expression of orexin receptors 1 and 2 in the rat brain. *J. Comp. Neurol.* 2001; 435(1): 6–25, doi: 10.1002/cne.1190.
22. Trivedi P., Yu H., MacNeil D.J., Van der Ploeg L.H., Guan X.M. Distribution of orexin receptor mRNA in the rat brain. *FEBS Lett.* 1998; 438(1–2): 71–75, doi: 10.1016/S0014-5793(98)01266-6.
23. Arias-Carrion O., Ortega-Robles E., de Celis-Alonso B., Palasz A., Mendez-Rojas M.A., Salas-Pacheco J. et al. Depletion of hypocretin/orexin neurons increases cell proliferation in the adult subventricular zone. *CNS Neurol. Disord. Drug Targets* 2018; 17(2): 106–112, doi: 10.2174/1871527317666180314115623.
24. Barson J.R., Leibowitz S.F. Orexin/hypocretin system: role in food and drug overconsumption. *Int. Rev. Neurobiol.* 2017; 136: 199–237, doi: 10.1016/bs.im.2017.06.006.
25. Nevárez N., de Lecea L. Recent advances in understanding the roles of hypocretin/orexin in arousal, affect, and motivation. *F1000Res.* 2018; 7: F1000 Faculty Rev-1421, doi: 10.12688/f1000research.15097.1.
26. Sargin D. The role of the orexin system in stress response. *Neuropharmacology* 2019; 154: 68–78, doi: 10.1016/j.neuropharm.2018.09.034.
27. Haghighparast A., Fatahi Z., Arezoomandan R., Karimi S., Taslimi Z., Zarabian S. Functional roles of orexin/hypocretin receptors in reward circuit. *Prog. Brain Res.* 2017; 235: 139–154, doi: 10.1016/bs.pbr.2017.08.005.
28. Wang C., Wang Q., Ji B., Pan Y., Xu C., Cheng B. et al. The orexin/receptor system: molecular mechanism and therapeutic potential for neurological diseases. *Front. Mol. Neurosci.* 2018; 11: 220, doi: 10.3389/fnmol.2018.00220.
29. Chien Y.L., Liu C.M., Shan J.C., Lee H.J., Hsieh M.H., Hwu H.G. et al. Elevated plasma orexin A levels in a subgroup of patients with schizophrenia associated with fewer negative and disorganized symptoms. *Psychoneuroendocrinology* 2015; 53: 1–9, doi: 10.1016/j.psyneuen.2014.12.012.
30. Lu X.Y., Bagnol D., Burke S., Akil H., Watson S.J. Differential distribution and regulation of OX1 and OX2 orexin/hypocretin receptor messenger RNA in the brain upon fasting. *Horm. Behav.* 2000; 37(4): 335–344, doi: 10.1006/hbeh.2000.1584.
31. Alizamini M.M., Kavianpour M., Karimi-Haghighi S., Fatahi Z., Haghighparast A. Intra-hippocampal administration of orexin receptor antagonists dose-dependently attenuates reinstatement of morphine seeking behavior in extinguished rats. *Peptides* 2018; 110: 40–46, doi: 10.1016/j.peptides.2018.10.011.
32. Mavanji V., Butterick T.A., Duffy C.M., Nixon J.P., Billington C.J., Kotz C.M. Orexin/hypocretin treatment restores hippocampal-dependent memory in orexin-deficient mice. *Neurobiol. Learn. Mem.* 2017; 146: 21–30, doi: 10.1016/j.nlm.2017.10.014.
33. Sadeghi B., Ezzatpanah S., Haghighparast A. Effects of dorsal hippocampal orexin-2 receptor antagonism on the acquisition, expression, and extinction of morphine-induced place preference in rats. *Psychopharmacology* 2016; 233(12): 2329–2341, doi: 10.1007/s00213-016-4280-3.
34. Akbari E., Naghdi N., Motamedi F. Functional inactivation of orexin 1 receptors in CA1 region impairs acquisition, consolidation and retrieval in Morris water maze task. *Behav. Brain Res.* 2006; 173(1): 47–52, doi: 10.1016/j.bbr.2006.05.028.
35. Bahramzadeh Zoeram S., Elahdadi Salmani M., Lashkarbolouki T., Goudarzi I. Hippocampal orexin receptor blocking prevented the stress-induced social learning and memory deficits. *Neurobiol. Learn. Mem.* 2019; 157: 12–23, doi: 10.1016/j.nlm.2018.11.009.
36. Edalat P., Kavianpour M., Zarabian S., Haghighparast A. Role of orexin-1 and orexin-2 receptors in the CA1 region of hippocampus in the forced swim stress- and food deprivation-induced reinstatement of morphine seeking behaviors in rats. *Brain Res. Bull.* 2018; 142: 25–32, doi: 10.1016/j.brainresbull.2018.06.016.
37. Pourreza P., Babapour V., Haghighparast A. Role of dorsal hippocampal orexin-1 receptors in modulation of antinociception induced by chemical stimulation of the lateral hypothalamus. *Physiol. Behav.* 2018; 185: 79–86, doi: 10.1016/j.physbeh.2017.12.036.
38. Brojeni M.S., Rashvand M., Haghighparast A. Role of orexin receptors within the dentate gyrus of the hippocampus in antinociception induced by chemical stimulation of the lateral hypothalamus in the tail-flick test as a model



- of acute pain in rats. *Physiol. Behav.* 2019; 209: 112595, doi: 10.1016/j.physbeh.2019.112595.
39. Chieffi S., Carotenuto M., Monda V., Valenzano A., Villano I., Precenzano F. et al. Orexin system: the key for a healthy life. *Front. Physiol.* 2017; 8: 357, doi: 10.3389/fphys.2017.00357.
40. Li S.B., Jones J.R., de Lecea L. Hypocretins, neural systems, physiology, and psychiatric disorders. *Curr. Psychiatry Rep.* 2016; 18(1): 7, doi: 10.1007/s11920-015-0639-0.
41. Wayne M.J., Armstrong D.L., Phelix C.F., Oomura Y. Orexin-A (hypocretin-1) and leptin enhance LTP in the dentate gyrus of rats in vivo. *Peptides* 2004; 25(6): 991–996, doi: 10.1016/j.peptides.2004.03.018.
42. Yang L., Zou B., Xiong X., Pascual C., Xie J., Malik A. et al. Hypocretin/orexin neurons contribute to hippocampus-dependent social memory and synaptic plasticity in mice. *J. Neurosci.* 2013; 33(12): 5275–5284, doi: 10.1523/JNEUROSCI.3200-12.2013.
43. Ito N., Yabe T., Gamo Y., Nagai T., Oikawa T., Yamada H. et al. I.c.v. administration of orexin-A induces an antidepressive-like effect through hippocampal cell proliferation. *Neuroscience* 2008; 157(4): 720–732, doi: 10.1016/j.neuroscience.2008.09.042.
44. Arendt D.H., Ronan P.J., Oliver K.D., Callahan L.B., Summers T.R., Summers C.H. Depressive behavior and activation of the orexin/hypocretin system. *Behav. Neurosci.* 2013; 127(1): 86–94, doi: 10.1037/a0031442.
45. Zhao X., Zhang R.x., Tang S., Ren Y.y., Yang W.x., Liu X.m. et al. Orexin-A-induced ERK1/2 activation reverses impaired spatial learning and memory in pentylenetetrazol-kindled rats via OX1R-mediated hippocampal neurogenesis. *Peptides* 2014; 54: 140–147, doi: 10.1016/j.peptides.2013.11.019.
46. Samokhina E., Samokhin A. Neuropathological profile of the pentylenetetrazol (PTZ) kindling model. *Int. J. Neurosci.* 2018; 128(11): 1086–1096, doi: 10.1080/00207454.2018.1481064.
47. Sousa-Ferreira L., Alvaro A.R., Aveleira C., Santana M., Brandão I., Kügler S. et al. Proliferative hypothalamic neurospheres express NPY, AGRP, POMC, CART and orexin-A and differentiate to functional neurons. *PLoS One* 2011; 6(5): e19745, doi: 10.1371/journal.pone.0019745.
48. Recabal A., Caprile T., García-Robles M.L.A. Hypothalamic neurogenesis as an adaptive metabolic mechanism. *Front. Neurosci.* 2017; 11: 190, doi: 10.3389/fnins.2017.00190.
49. Rojczyk-Golebiewska E., Pałasz A., Wiaderekiewicz R. Hypothalamic subependymal niche: a novel site of the adult neurogenesis. *Cell. Mol. Neurobiol.* 2014; 34(5): 631–642, doi: 10.1007/s10571-014-0058-5.
50. Batailler M., Droguerre M., Baroncini M., Fontaine C., Prevot V., Migaud M. DCX-expressing cells in the vicinity of the hypothalamic neurogenic niche: a comparative study between mouse, sheep, and human tissues. *J. Comp. Neurol.* 2014; 522(8): 1966–1985, doi: 10.1002/cne.23514.
51. Ebling F.J. Hypothalamic control of seasonal changes in food intake and body weight. *Front. Neuroendocrinol.* 2015; 37: 97–107, doi: 10.1016/j.yfrne.2014.10.003.
52. Samms R.J., Lewis J.E., Lory A., Fowler M.J., Cooper S., Warner A. et al. Antibody-mediated inhibition of the FGFR1c isoform induces a catabolic lean state in Siberian hamsters. *Curr. Biol.* 2015; 25(22): 2997–3003, doi: 10.1016/j.cub.2015.10.010.
53. Rizzoti K., Lovell-Badge R. Pivotal role of median eminence tanycytes for hypothalamic function and neurogenesis. *Mol. Cell. Endocrinol.* 2017; 445: 7–13, doi: 10.1016/j.mce.2016.08.020.
54. Lee D.A., Bedont J.L., Pak T., Wang H., Song J., Miranda-Angulo A. et al. Tanycytes of the hypothalamic median eminence form a diet-responsive neurogenic niche. *Nat. Neurosci.* 2012; 15(5): 700–702, doi: 10.1038/nn.3079.
55. Lee D.A., Yoo S., Pak T., Salvatierra J., Velarde E., Aja S. et al. Dietary and sex-specific factors regulate hypothalamic neurogenesis in young adult mice. *Front. Neurosci.* 2014; 8: 157, doi: 10.3389/fnins.2014.00157.
56. Duque A., Spector R. A balanced evaluation of the evidence for adult neurogenesis in humans: implication for neuropsychiatric disorders. *Brain Struct. Funct.* 2019; 224(7): 2281–2295, doi: 10.1007/s00429-019-01917-6.